



This is an open access article distributed in accordance with the Creative Commons Attribution (CC BY 4.0) license: <https://creativecommons.org/licenses/by/4.0/> which permits any use, Share — copy and redistribute the material in any medium or format, Adapt — remix, transform, and build upon the material for any purpose, as long as the authors and the original source are properly cited. © The Author(s) 2025

Impending role of biomarkers of oxidative stress in myocardial infarction: a comparative study from local population

Arif Malik^{1,2*}, Anoosh Qayum³, Jehanzaib Islam⁴, Ayesha Zahid¹, Haleema Saadia¹, Gul Zaib¹

This article has been retracted.

ABSTRACT

Background and Objective: Myocardial infarction (MI) causes morbidity and mortality worldwide and is linked to oxidative stress and metabolic markers. This study examines how oxidative stress markers predict myocardial infarction and effect its course, thus having potential for use as markers for early MI diagnosis.

Method: This cross-sectional comparative study was conducted at the University of Lahore, Institute of Molecular Biology and Biotechnology, on 405 clinically proven MI patients with age-matched controls of 20-90 years. Biochemical marker concentrations were determined using standardized ELISA kits on blood samples containing vascular smooth muscle cells. The study examined markers such as lipid peroxidation (DT, MDA, THB, 8-OHdG, 4-HNE, NT, and IsoP-F2 α) and antioxidants (SOD, Catalase).

Results: All variables demonstrated a statistically significant difference between MI patients and controls. Serum MDA levels in MI patients were higher than the control group (4.56 ± 1.09 versus 0.932 ± 0.056 nmol/ml, $p = 0.022$). In the MI group, serum oxidative stress levels (8-OHdG, 4-HNE, IsoP-F2 α) were greater than in the control group. Patients with MI had reduced SOD, CAT, GSH, Vitamin D, C, and E levels.

Conclusion: In conclusion, oxidative stress and antioxidant enzyme levels predict myocardial infarction outcomes. Low vitamin and antioxidant levels and high MDA, 4-HNE, and isoprostanes levels predispose to MI. The annotations show early patterns and treatment options for oxidative stress in cardiovascular patients.

Keywords: Myocardial infarction, oxidative stress, MDA, 4-HNE, predictive markers.

Received: 26 September 2024

Revised (1): 23 November 2024

Revised (2): 14 December 2024

Accepted: 09 January 2025

Correspondence to: Arif Malik

*School of Pain and Regenerative Medicine, The University of Lahore, Lahore, Pakistan and Faculty of Health Sciences, Equator University of Science and Technology, (EQUaT), Masaka, Uganda.

Email: arifuaf@yahoo.com

Full list of author information is available at the end of the article.

Authors' Details

Arif Malik^{1,2}, Anoosh Qayum³, Jehanzaib Islam⁴, Ayesha Zahid¹, Haleema Saadia¹, Gul Zaib¹

1. School of Pain and Regenerative Medicine, The University of Lahore, Lahore, Pakistan

2. Faculty of Health Sciences, Equator University of Science and Technology, (EQUaT), Masaka, Uganda

3. Department of Biochemistry, Ameer Ud Din Medical College, Postgraduate Medical Institute, Lahore, Pakistan

4. Department of Biological Sciences, Grand International University, Sialkot, Pakistan